



LASIK POSTOP

Hoopes Vision
11820 S. State Street
Draper, UT 84020
801.568.0200



ANTIBIOTIC:
Moxifloxacin/Ofloxacin
**EVERY 4 HOURS
FOR 1 WEEK**

Wait at least
5 minutes
between drops.
Only necessary
during waking
hours.

**ARTIFICIAL TEARS
(PRESERVATIVE FREE)**



STEROID:
Prednisolone Acetate
**EVERY 2 HOURS
UNTIL YOUR 1 DAY POSTOP VISIT
THEN
EVERY 4 HOURS
FOR 1 WEEK**

**1-2 Drops EVERY 15-30 MIN
THE DAY OF SURGERY
THEN
AT LEAST EVERY HOUR
FOR 1 MONTH**

* SHAKE BEFORE
EACH USE

AFTER YOUR SURGERY:


- A short nap is advised. Limit activity and phone used the day of surgery. Increase activity as your comfort and vision allows. Take your normal over-the-counter pain medication if needed.
- Your vision will be foggy for the first day.
- You may experience some tearing, burning, stinging, light sensitivity, redness, and foreign body sensation.
- Do not drive until after your 1-day post-op appointment. You may return to work and/or drive when you are confident with your vision and your comfort allows.
- If you notice increasing pain or a sudden decrease in vision – call us immediately.

ARTIFICIAL TEARS:

- FIRST MONTH – Use preservative free artificial tears in vials 1-2 drops **every hour**
- MONTH 2 & 3 - OK to switch to a bottled artificial tear and use every 2 to 3 hours.

OTHER IMPORTANT PRECAUTIONS:

- Wash and dry your hands before putting in your eye drops. Be careful not to touch the tip of the bottle to any part of your eye. You may notice an unusual taste after using some drops.
- Your vision may fluctuate for the first several weeks.
- You may experience halos and glare around lights at night for the first couple weeks.
- No eye makeup for 3 - 5 days. New mascara is recommended after the procedure.
- Avoid getting soap or water in your eyes while showering for the first week.
- Wear your shields while sleeping or napping for the first week.
- You should protect your eyes from bright sun and wind for 1 month.
- No swimming or hot tubs for 2 weeks. No high-impact water sports for 3 months.
- Do not rub or bump your eyes for at least 2 months after surgery. It is a bad habit to rub your eyes. We suggest you avoid this habit altogether.



Preservative Free
Artificial Tears (vials)
Month 1



OK to use Bottle
Artificial Tears
when used less
frequently

Hoopes Vision Phone Number: 801-568-0200

Benjamin Buckner, MD

Patient Signature: _____

Date: _____

Staff Signature: _____

Tomorrow's Appointment Time: